



A healthy workplace is good for business

HEALTH EFFECTS OF SMOKING

Tobacco smoking costs Australia over \$30 billion a year in health, business and social costs. The costs to business alone are over \$5.7 billion.¹ Smoking is the single greatest cause of preventable death in the developed world.²

In 2011-2012, the prevalence of smoking amongst Tasmanian adults was 23.2%, significantly above the national average of 18.1%. Smoking prevalence was higher amongst males (28%) compared to females (18.6%).³

The best thing smokers can do for their health is to quit smoking.

Health effects for smokers

The health effects of cigarette smoke have been well researched for several decades. Smokers have more health issues and a lower life expectancy than the general population. Smoking causes a range of serious health problems, including cancer, heart disease, stroke, asthma, emphysema, vascular disease and damage to most body organs.

Smokers' health problems can impact on their work performance and affect productivity through more frequent sickness-related absences and multiple 'smoke breaks' while at work.⁴

Despite common thinking that smoking is a stress reliever, research has found that it actually increases psychological distress while providing no medicinal benefits.⁵





Health effects for non-smokers

Non-smokers can be affected by inhaling environmental tobacco smoke released from a burning cigarette or smoke that is exhaled by a smoker.

Environmental tobacco smoke is the mixture of chemicals and toxins released whenever someone smokes a cigarette. Many of the 4000+ chemicals in tobacco smoke are poisonous, and at least 69 are known cancer-causing agents.⁶ 'Passive smoking' occurs when non-smokers inhale environmental tobacco smoke causing non-smokers to breathe the same dangerous substances as smokers.

Some of the immediate effects of passive smoking are eye irritation, headache, cough and sore throat.^{7,8,9} Passive smoking can also lead to more serious health issues, similar to those of smokers, including lung cancer, heart disease and respiratory problems.^{7,8,9} There are health risks from being exposed to environmental tobacco smoke in both indoor and outdoor areas.

A report by the US Surgeon General has concluded that there is no safe level of exposure to environmental tobacco smoke.⁷

References

1. Collins DJ, Lapsley, HM 2008, *The costs of tobacco, alcohol, and illicit drugs to Australian society in 2004-2005*, Commonwealth of Australia, p. xi and table 33, p. 64.
2. World Health Organisation 2002, *Smoking statistics*, http://www.wpro.who.int/mediacentre/factsheets/fs_20020528/en/index.html
3. Australian Bureau of Statistics 2012, *Australian Health Survey: 2011-2012 - Tasmania*, Australian Government, Canberra.
4. Halpern, MT, Shikiar, R, Rentz, AM 2001, 'Impact of smoking status on workplace absenteeism and productivity', *Tobacco Control*, vol. 10, pp. 233-238.
5. Parrott, AC, 2003 'Cigarette-derived nicotine is not a medicine', *World Journal of Biological Psychiatry*, vol. 4, pp. 49-55.
6. Quit Victoria 2006, *Quit because you can (booklet)*, Victoria.
7. US Department of Health and Human Services 2006, *The health consequences of involuntary exposure to tobacco smoke: a report of the Surgeon General*, US Department of Health and Human Services, Centres for Disease Control and Prevention, Coordinating Centre for Health Promotion, National Centre for Chronic Disease Prevention and Health Promotion and Office on Smoking and Health, Atlanta.
8. California Environmental Protection Agency, Office of Environmental Health Hazard Assessment 2005, *Proposed identification of environmental tobacco smoke as a toxic air contaminant*, Appendix 3, pp. 10-12.
9. World Health Organisation: International Agency for Research on Cancer (IARC) 2004, *Tobacco Smoke and Involuntary Smoking*, Monograph 83, IARC, Lyon.